

## Mid-infrared predictions of cheese yield from bovine milk

Vanlierde A.<sup>1,2</sup>, Soyeurt H.<sup>1,3</sup>, Anceau C.<sup>2</sup>, Vanden Bossche S.<sup>2</sup>, Dehareng F.<sup>4</sup>, Dardenne P.<sup>4</sup>, Gengler N.<sup>1,3</sup>, Sindic M.<sup>2</sup> and Colinet F.G.<sup>1,\*</sup>

1 Animal Science Unit, Gemblooux Agro-Bio Tech, University of Liège, Belgium  
 2 Analysis, Quality and Risk Unit, Gemblooux Agro-Bio Tech, University of Liège, Belgium  
 3 National Fund for Scientific Research, Belgium  
 4 Valorisation of Agricultural Products Department, Walloon Agricultural Research Center, Belgium  
 \* Presenting author's email: [Frederic.Colinet@ulg.ac.be](mailto:Frederic.Colinet@ulg.ac.be)

62<sup>nd</sup> EAAP Annual Meeting, Stavanger, Norway, August 29<sup>th</sup> - September 1<sup>st</sup> 2011

## Context

- Cheese manufacture
  - Economical importance
  - Empirical & theoretical formula for cheese yield (CY)
    - ✓ Based on some factors:
      - Milk fat content
      - Milk protein content
      - Milk casein content
      - Moisture
      - Salt
      - ...

62<sup>nd</sup> EAAP Annual Meeting, Stavanger, Norway, August 29<sup>th</sup> - September 1<sup>st</sup> 2011

## Context

- Cheese yield
  - Influence of animal selection on milk component
    - ➔ also on milk processability
  - Interest for studying CY at large scale

But the CY measurement on a large number of individual samples is

  - ✓ Time consuming
  - ✓ Skilled staff

**Request for a CY prediction without the need of the prior determination of milk composition**

62<sup>nd</sup> EAAP Annual Meeting, Stavanger, Norway, August 29<sup>th</sup> - September 1<sup>st</sup> 2011

## Objective

- To investigate the potential use of Mid Infrared (MIR) spectrometry in order to predict cheese yield

**Why MIR spectrometry ?**

- ➔ Small quantity of milk and fast method
- ➔ Already implemented in milk labs to measure the contents of fat, protein, lactose, and urea in milk

62<sup>nd</sup> EAAP Annual Meeting, Stavanger, Norway, August 29<sup>th</sup> - September 1<sup>st</sup> 2011

## Material and methods

- Sampling
  - Walloon Region of Belgium
  - Variability of spectra: several criteria
    - ✓ Milk sampling: individual or bulk milk
    - ✓ Breed: Dual Purpose Belgian Blue, (Red) Holstein, Montbéliarde and Jersey
    - ✓ Time of sampling: morning milking, evening milking or mix of 50 % morning & 50 % evening milk samples

➔ 157 samples collected (October 2009 – May 2010)

62<sup>nd</sup> EAAP Annual Meeting, Stavanger, Norway, August 29<sup>th</sup> - September 1<sup>st</sup> 2011

## Material and methods

- Analysis
  - Milk Lab (Comité du Lait, Battice, Belgium)
    - ✓ MIR Foss MilkoScanFT6000 spectrometer
    - ✓ Analysed traits: fat, protein, lactose, somatic cell count (SCC) and pH
    - ✓ SCC ➔ Somatic Cell Score (SCS)

62<sup>nd</sup> EAAP Annual Meeting, Stavanger, Norway, August 29<sup>th</sup> - September 1<sup>st</sup> 2011

## Material and methods

### ■ Analysis

- Individual Laboratory Cheese Yield (ILCY) were determined according to Hurtaud *et al.*, 1995 (Ann. Zootech. 44, 385-398)
  - ✓ g dry coagulum / 100 g milk dry matter
  - ✓ 2 samples discarded due to poor coagulation

62<sup>nd</sup> EAAP Annual Meeting, Stavanger, Norway, August 29<sup>th</sup> - September 1<sup>st</sup> 2011

7

## Material and methods

### ■ MIR-Chemometric methods

- WINISI III software
- Partial Least Square (PLS) regressions
- Use (or not) of a 1<sup>st</sup> derivative pretreatment (1<sup>st</sup>Der)
  - ✓ Permits to correct the baseline drift
- Detection of outlier spectrum
  - ✓ Based on PCA and Mahalanobis distance
- Use (or not) of a repeatability file (RepFile)
  - ✓ Spectra from the same samples analysis on different spectrometers

62<sup>nd</sup> EAAP Annual Meeting, Stavanger, Norway, August 29<sup>th</sup> - September 1<sup>st</sup> 2011

8

## Material and methods

### ■ MIR-Chemometric methods

- Statistical parameters
  - ✓ Mean and standard deviation (SD)
  - ✓ Standard error of calibration (SEC)
  - ✓ Calibration coefficient of determination ( $R^2_c$ )
- Internal validation by leave one out cross-validation

62<sup>nd</sup> EAAP Annual Meeting, Stavanger, Norway, August 29<sup>th</sup> - September 1<sup>st</sup> 2011

9

## Material and methods

### ■ MIR-Chemometric methods

- Statistical parameters to assess the accuracy
  - ✓ Standard error of cross-validation (SECV)
  - ✓ Cross-validation coefficient of determination ( $R^2_{cv}$ )
- Efficiency of calibration
  - ✓ RPD = SD / SECV

62<sup>nd</sup> EAAP Annual Meeting, Stavanger, Norway, August 29<sup>th</sup> - September 1<sup>st</sup> 2011

10

## Material and methods

### ■ MIR-Chemometric methods

- T-outlier test
  - ✓ Compare observed and predicted values
  - ✓ Samples with T outlier value > 2.5 were discarded
  - ✓ Maximum 5 tests performed

62<sup>nd</sup> EAAP Annual Meeting, Stavanger, Norway, August 29<sup>th</sup> - September 1<sup>st</sup> 2011

11

## Material and methods

### ■ MIR-Chemometric methods

- 4 methods tested
  - ✓ PLS
  - ✓ PLS + RepFile
  - ✓ PLS + 1<sup>st</sup>Der
  - ✓ PLS + 1<sup>st</sup>Der + RepFile

62<sup>nd</sup> EAAP Annual Meeting, Stavanger, Norway, August 29<sup>th</sup> - September 1<sup>st</sup> 2011

12

## Results

### Characterization of the 155 samples

Trait	Mean	SD
Fat (%)	3.81	1.03
Protein (%)	3.57	0.50
Lactose (g/100 mL)	4.83	0.27
DM (%)	12.82	2.57
SCS	3.18	1.96
pH	6.67	0.07
ILCY (g dry coagulum / 100 g milk DM)	62.8	12.6

DM = Dry Matter; SCS = Somatic Cell Score;  
ILCY = Individual Laboratory Cheese Yield

62<sup>nd</sup> EAAP Annual Meeting, Stavanger, Norway, August 29<sup>th</sup> - September 1<sup>st</sup> 2011

13

## Results

### Characterization of the 155 samples

Trait	Mean	SD
Fat (%)	3.81	1.03
Protein (%)	3.57	0.50
Lactose (g/100 mL)	4.83	0.27
DM (%)	12.82	2.57
SCS	3.18	1.96
pH	6.67	0.07
ILCY (g dry coagulum / 100 g milk DM)	62.8	12.6

Coefficient of Variation = 20 %

DM = Dry Matter; SCS = Somatic Cell Score;  
ILCY = Individual Laboratory Cheese Yield

62<sup>nd</sup> EAAP Annual Meeting, Stavanger, Norway, August 29<sup>th</sup> - September 1<sup>st</sup> 2011

14

## Results

### Observed correlations among milk components

	Fat	Protein	Lactose	DM	SCS	pH
ILCY	0.50***	0.47***	-0.20*	0.24**	0.31***	0.07 <sup>NS</sup>

ILCY = Individual Laboratory Cheese Yield; DM = Dry Matter;  
SCS = Somatic Cell Score; \* = *P*-value < 0.05;  
\*\* = *P*-value < 0.01; \*\*\* = *P*-value < 0.001; NS = non significant

62<sup>nd</sup> EAAP Annual Meeting, Stavanger, Norway, August 29<sup>th</sup> - September 1<sup>st</sup> 2011

15

## Results

### Calibration equations

Methods	No. outlier spectra	T-outlier test	
		Iteration	No. outlier
PLS	3	5	15
PLS + RepFile	3	5	23
PLS + 1 <sup>st</sup> Der	1	4	12
PLS + 1 <sup>st</sup> Der + RepFile	1	3	22

62<sup>nd</sup> EAAP Annual Meeting, Stavanger, Norway, August 29<sup>th</sup> - September 1<sup>st</sup> 2011

16

## Results

Similar  $R^2_c$  and  $R^2_{cv}$   
Low RPD

### Calibration equations

	PLS	PLS RepFile	PLS 1 <sup>st</sup> Der	PLS 1 <sup>st</sup> Der RepFile
N	137	129	142	132
SD	12.00	11.56	11.97	11.81
No. of factor	6	5	9	9
$R^2_c$	0.72	0.66	0.76	0.76
$R^2_{cv}$	0.68	0.62	0.66	0.68
RPD	1.75	1.61	1.70	1.76

SD = Standard Deviation;  $R^2_c$  = Calibration Coefficient of determination;  
 $R^2_{cv}$  = Cross-Validation Coefficient of determination;  
RPD = ratio of SD to the standard error of cross-validation


62<sup>nd</sup> EAAP Annual Meeting, Stavanger, Norway, August 29<sup>th</sup> - September 1<sup>st</sup> 2011

17

## Results

### Developed equation using PLS + 1<sup>st</sup>Der + RepFile

#### Scatter plot of Observed vs. Predicted ILCY




62<sup>nd</sup> EAAP Annual Meeting, Stavanger, Norway, August 29<sup>th</sup> - September 1<sup>st</sup> 2011

18

## Results

- Developed equation using PLS + 1<sup>st</sup>Der + RepFile
- Scatter plot of Residuals vs. Observed ILCY

62<sup>nd</sup> EAAP Annual Meeting, Stavanger, Norway, August 29<sup>th</sup> - September 1<sup>st</sup> 2011

19

## Conclusions

- $R^2_C$  and  $R^2_{CV}$  of developed equations
  - Interesting coefficients of determination
  - But  $< 0.80$
- RPD  $< 2$

➔ First results are promising for the prediction of an indicator for ILCY from MIR spectrum  
 ➔ Dataset should be enriched with new samples in order to better control the variability

62<sup>nd</sup> EAAP Annual Meeting, Stavanger, Norway, August 29<sup>th</sup> - September 1<sup>st</sup> 2011

20

## Perspectives

- Improvement with additional samples
- Use of the developed equation
  - Walloon Database: 1,800,000 spectra
  - Study of ILCY variability in the Walloon dairy cattle
    - ✓ Detection of potential effects of breed, season, DIM...
  - Development of a genetic evaluation

62<sup>nd</sup> EAAP Annual Meeting, Stavanger, Norway, August 29<sup>th</sup> - September 1<sup>st</sup> 2011

21

## Thank you for your attention



Corresponding author's e-mail: Frederic.Colinet@ulg.ac.be

- Acknowledgments for financial support
  - European Commission (ERDF) and SPW – DGO3 through projects Interreg IVa BlueSel and D31-1255/S1
  - National Fund for Scientific Research through travel grant for attending EAAP annual meeting
- Acknowledgments for collaboration
  - Milk Committee of Battice (Belgium)
  - Walloon Breeding Association (AWE asbl)
  - Walloon dairy breeders

62<sup>nd</sup> EAAP Annual Meeting, Stavanger, Norway, August 29<sup>th</sup> - September 1<sup>st</sup> 2011